

# Enhanced Dimer Relaxation in an Atomic/Molecular BEC

Eric Braaten\*

Department of Physics, The Ohio State University, Columbus, OH 43210, USA

H.-W. Hammer†

Helmholtz-Institut für Strahlen- und Kernphysik (Abt. Theorie), Universität Bonn, 53115 Bonn, Germany‡

(Dated: June, 2004)

We derive a universal formula for the rate constant  $\beta$  for relaxation of a shallow dimer into deeply-bound diatomic molecules in the case of atoms with a large scattering length  $a$ . We show that  $\beta$  is determined by  $a$  and by two 3-body parameters that also determine the binding energies and widths of Efimov states. The rate constant  $\beta$  scales like  $\hbar a/m$  near the resonance, but the coefficient is a periodic function of  $\ln(a)$  that may have resonant enhancement at values of  $a$  that differ by multiples of 22.7.

PACS numbers: 03.75.Nt, 34.50.-s, 21.45.+v

Keywords: Dimer relaxation, universality, large scattering length

Using Feshbach resonances [1], it is possible to study cold atoms with a variable scattering length  $a$  that can be made large enough that the atoms have strong resonant interactions. Surprising phenomena have been observed in experiments on cold atoms with large scattering lengths. Experiments with a  $^{23}\text{Na}$  BEC [2], ultracold  $^{85}\text{Rb}$  atoms [3], a  $^{87}\text{Rb}$  BEC [4], and ultracold  $^{133}\text{Cs}$  atoms [5] have revealed large losses of atoms as the Feshbach resonance is approached. In experiments with a  $^{85}\text{Rb}$  BEC in which  $a$  was increased briefly to large positive values, a burst of energetic atoms was observed along with missing atoms [6]. More recently, atom-molecule coherence was demonstrated [7], suggesting the coexistence of a condensate of shallow diatomic molecules with an atom condensate. Few-body processes play an important role in these phenomena and it is important to understand their behavior in the limit of large  $a$ .

A Feshbach resonance allows  $|a|$  to be made large compared to the typical low-energy length scale  $\ell = (mC_6/\hbar^2)^{1/4}$  set by the atom mass  $m$  and the van der Waals coefficient  $C_6$ . If  $|a| \gg \ell$ , the few-body problem exhibits universal properties that are insensitive to the details of the interactions responsible for the large scattering length. A simple example in the 2-body sector for  $a > 0$  is the existence of a shallow molecule (the *dimer*) with binding energy  $|E_2| = \hbar^2/(ma^2)$ . A particularly remarkable example in the 3-body sector is the existence of shallow 3-body bound states (*Efimov states*) with universal properties [8]. As  $|a|$  increases, shallower Efimov states appear below the scattering threshold at values of  $a$  that differ by multiples of  $e^{\pi/s_0} \approx 22.7$ , where  $s_0 \approx 1.00624$ . The Efimov states can cause dramatic dependence of scattering observables on  $a$  and on the energy. The 3-body scattering observables are determined

by  $a$  and by a 3-body parameter that also determines the Efimov spectrum. Simple universal expressions have been derived for the *atom-dimer scattering length* [8, 9] and for the rate constant for *3-body recombination* into the shallow dimer [10, 11, 12]. They are valid for all atoms with  $|a| \gg \ell$  that have no 2-body bound states if  $a < 0$  and only the shallow dimer if  $a > 0$ .

Unfortunately, heavy alkali atoms, such as Rb and Na, form many *deeply-bound diatomic molecules*. A simple consequence is that Efimov states are resonances rather than sharp states, because they can decay into a deep molecule and an energetic atom. The widths of Efimov resonances have been studied in Refs. [13] using explicit model potentials. The deep molecules also affect other 3-body observables. In particular, the universal expressions for the atom-dimer scattering length and for the 3-body recombination rate constant must be modified.

The existence of deep molecules opens up additional channels for the loss of cold atoms from an ultracold gas or a BEC. One such channel is 3-body recombination into deep molecules, which has been studied previously in Refs. [11, 14]. Another loss mechanism for a cold gas containing both atoms and dimers is *dimer relaxation*, in which the collision of an atom and a shallow dimer produces a deep molecule and a recoiling atom. If the momenta of the incoming atom and dimer are sufficiently small, the relaxation rate reduces to a constant. The *relaxation event rate constant*  $\beta$  is defined so that the number of relaxation events per time and per volume in a cold gas (or BEC) of atoms and dimers with number densities  $n_A$  and  $n_D$  is  $\beta n_A n_D$ . Because of the large energy released, this process results in the loss of a low-energy atom and a shallow dimer:

$$\frac{dn_A}{dt} = \frac{dn_D}{dt} = -\beta n_A n_D. \quad (1)$$

The loss of atoms from a BEC through dimer relaxation has been considered previously in Refs. [15]. The authors assumed that the relaxation rate constant  $\beta$  is independent of  $a$ . They found qualitative agreement with the slow-sweep experiments in [2] for  $\beta \approx 10^{-10} \text{ cm}^3/\text{s}$ .

\*Electronic address: braaten@mps.ohio-state.edu

†Electronic address: hammer@phys.washington.edu

‡Present Address: Institute for Nuclear Theory, University of Washington, Seattle, WA 98195, USA

In this paper, we derive a universal expression for the dimer relaxation rate constant  $\beta$  in the limit of large  $a$  using hyperspherical methods combined with previous results from an effective field theory. We show that  $\beta$  scales like  $\hbar a/m$ , but with a coefficient that is a periodic function of  $\ln(a)$  with period  $\pi/s_0 \approx \ln(22.7)$ . The coefficient of  $\hbar a/m$  also depends on two 3-body parameters that determine the binding energies and widths of the Efimov resonances. It can have resonant enhancement at those values of  $a$  for which there is an Efimov resonance near the atom-dimer threshold.

In contrast to other treatments of Feshbach resonances, we do not attempt to describe the details of the physics in the short-distance region. Such details include the closed channel of the Feshbach resonance and the decomposition of the dimer relaxation rate into contributions from individual deep 2-body bound states. A detailed description of the short-distance physics is not necessary for calculating the inclusive dimer relaxation rate. Since the total probability is conserved, one can describe the inclusive rate by taking advantage of unitarity.

We first consider the case without deep dimers. Any probability that flows to the short-distance region where the atoms are coupled to the closed channel must be reflected back to the long-distance region in the form of low-momentum states. A detailed knowledge of the short-distance dynamics of the Feshbach resonance which creates the large scattering length is not required. One can eliminate the closed channel in favor of the scattering length  $a$  and the 3-body parameter  $\Lambda_*$  discussed below. This simplification is possible because the resolution scale of the low-momentum atoms is given by the scattering length  $a$  which is much larger than the range of the potential for the closed channel. As a consequence, low-energy atoms are only able to feel the large scattering length  $a$  (and the three-body parameter  $\Lambda_*$  if three- and higher-body observables are concerned) but not to probe the mechanism responsible for it.

If deep dimers are present, some of the probability that flows to the short-distance region will emerge in the form of high-momentum states consisting of an atom and a deep dimer. However, if one is interested in the inclusive relaxation rate only, it is irrelevant how exactly the probability is lost from the low-momentum states. All one needs to know is how much of the probability is lost. In the case of atoms with a large scattering length, this can be parametrized by one number  $\eta_*$ . The reason for this is again that the resolution scale of the low-momentum atoms is given by the scattering length  $a$  which is much larger than the size of the deep dimer. Therefore the low-momentum atoms cannot resolve the details of the physics at short distances comparable to the size of the deep dimer.

A similar approach was previously used to describe the effect of deep 2-body bound states on Efimov states [16] or the three-body recombination rate [14]. It has also been applied to many other problems in physics. In the mean-field equations for Bose-Einstein condensates,

atom loss processes are often taken into account through imaginary parts of coefficients in the mean-field Hamiltonian [17]. If the loss processes involve momenta large compared to those of atoms in the condensate, this prescription can be rigorously justified. A similar method is used in nuclear and particle physics to describe effects of the annihilation of particles and their antiparticles [18]. The dynamics of the short-distance annihilation process may be very complicated, but its effects on low-momentum particles can be described by local operators in an effective Hamiltonian and all one needs to know is the imaginary parts of the coefficients of those terms.

In the remainder of the paper, we will make these arguments more quantitative and use them to derive a universal expression for the dimer relaxation rate constant  $\beta$  in the limit of large  $a$ . We begin by considering the equation for the hyperradial wavefunction  $f(R)$  for the lowest adiabatic potential in the hyperspherical representation of the 3-body problem [19]. In the region  $\ell \ll R \ll a$ , the other adiabatic potentials are repulsive and the corresponding wavefunctions decrease exponentially at  $R \ll a$ . The coupling to those channels can therefore be neglected and the equation for  $f(R)$  reduces to

$$-\frac{\hbar^2}{2m} \left[ \frac{\partial^2}{\partial R^2} + \frac{s_0^2 + 1/4}{R^2} \right] f(R) = E f(R). \quad (2)$$

This looks like the Schrödinger equation for a particle in a one-dimensional, scale-invariant  $1/R^2$  potential. For low energies  $|E| \sim \hbar^2/ma^2$ , the energy eigenvalue in Eq. (2) can be neglected and the most general solution is

$$f(R) = \sqrt{HR} \left[ A e^{i s_0 \ln(HR)} + B e^{-i s_0 \ln(HR)} \right], \quad (3)$$

where  $H$  has dimensions of wave number but is otherwise arbitrary. Equation (3) is simply the sum of outgoing and incoming hyperradial waves. At shorter distances  $R \sim \ell$  and longer distances  $R \sim a$ , the wave function becomes very complicated, but it is constrained by unitarity.

Following Efimov [8], we first exploit unitarity in the long-distance region  $R \sim a$ . If we identify the asymptotic states at distances  $R \ll a$  and  $R \gg a$ , the evolution of the wave function through the region  $R \sim a$  can be described by a unitary matrix  $s$  (see the illustration in Fig. 1). We denote the asymptotic states with probability flowing into and out of the region  $R \sim a$  by kets  $|i \text{ in}\rangle$  and  $|i \text{ out}\rangle$ . The probability can flow into this region either from  $\ell \ll R \ll a$  or from  $R \gg a$ . In the region  $\ell \ll R \ll a$ , the asymptotic states  $|1 \text{ in}\rangle$  and  $|1 \text{ out}\rangle$  are the outgoing and incoming hyperradial waves represented by the first and second terms in Eq. (3). For simplicity, we restrict ourselves to energies below the dimer-breakup threshold where 3-atom states are kinematically forbidden. In the region  $R \gg a$ , the asymptotic states  $|2 \text{ in}\rangle$  and  $|2 \text{ out}\rangle$  are incoming and outgoing atom-dimer scattering states. The amplitudes for incoming asymptotic states  $|j \text{ in}\rangle$  to evolve into outgoing asymptotic states  $|i \text{ out}\rangle$  form a unitary  $2 \times 2$  matrix  $s_{ij}$ . Time-reversal invariance implies that  $s$  is a symmetric matrix. Since  $s$  is




FIG. 1: The hyperspherical potential in the region  $l \ll R \ll a$ . The evolution of wavefunctions in the regions  $R \sim a$  and  $R \sim l$  is described by unitary matrices  $s$  and  $t$ .

dimensionless, it can depend on the interaction potential and the energy only through the variable  $ma^2E/\hbar^2$ .

Following Efimov, we next exploit unitarity in the short-distance region  $R \sim \ell$ . If there are no deep molecules, the probability in the incoming hyperradial wave must be totally reflected from the region  $R \sim \ell$  [8]. Thus the amplitudes of the incoming and outgoing hyperradial waves in (3) must be equal in magnitude:  $A = -Be^{2i\theta_*}$ . The angle  $\theta_*$  is determined by the logarithmic derivative  $f'(R)/f(R)$  in the region  $R \ll a$ . This implies that  $\theta_*$  must have the form  $\theta_* = s_0 \ln(Ca\Lambda_*)$ , where  $C$  is a dimensionless numerical constant and  $\Lambda_*$  is a parameter that is insensitive to variations in  $a$ . The evolution of the wave function in the region  $R \sim \ell$  can be described by a  $1 \times 1$  unitary matrix whose only entry is  $t_{11} = e^{2i\theta_*}$ . By combining the constraints from unitarity in the regions  $R \sim \ell$  and  $R \sim a$ , Efimov obtained an expression for the S-matrix element for atom-dimer scattering that he referred to as a radial law [8]:

$$S_{AD,AD} = s_{22} + s_{21}(1 - t_{11}s_{11})^{-1}t_{11}s_{12}. \quad (4)$$

Because  $s$  is dimensionless, it can only depend on the dimensionless variable  $ma^2E/\hbar^2$ . The atom-dimer scattering phase shift can therefore be obtained by calculating a few universal functions of  $ma^2E/\hbar^2$  [8].

We now consider the case with deeply-bound diatomic molecules. In this case, some of the probability in an incoming hyperradial wave that flows into the region  $R \sim \ell$  will emerge as high-energy scattering states of an atom and a deep molecule. The amplitude of the outgoing hyperradial wave in (3) must therefore be smaller than that of the incoming hyperradial wave:  $A = -Be^{2i\theta_* - 2\eta_*}$ . The corresponding element of the unitary matrix  $t$  is

$$t_{11} = \exp(2i\theta_* - 2\eta_*). \quad (5)$$

Inserting this expression into (4), we obtain the generalization of Efimov's radial law to the case in which there are deep molecules. If the universal expressions for the

S-matrix elements are known in the case  $\eta_* = 0$ , the effects of deep molecules can be deduced by the simple substitution  $\theta_* \rightarrow \theta_* + i\eta_*$  [16].

We proceed to derive the radial law for S-matrix elements involving the deep molecules. In the short-distance region  $R \sim \ell$ , the relevant asymptotic states are incoming and outgoing hyperradial waves  $|1 \text{ in}\rangle$  and  $|1 \text{ out}\rangle$  and high-energy atom-molecule scattering states which we denote by  $|X \text{ in}\rangle$  and  $|X \text{ out}\rangle$ . Note that the outgoing hyperradial wave is an incoming asymptotic state  $|1 \text{ in}\rangle$  as far as the region  $R \sim a$  is concerned and an outgoing asymptotic state  $|1 \text{ out}\rangle$  as far as the region  $R \sim \ell$  is concerned. The S-matrix element for a low-energy atom-dimer scattering state to evolve into a high-energy atom-molecule scattering state  $X$  is

$$S_{X,AD} = t_{X1}s_{12} + t_{X1}s_{11}(1 - t_{11}s_{11})^{-1}t_{11}s_{12}. \quad (6)$$

The first term on the right side of Eq. (6) describes transmission through both the regions  $\ell \ll R \ll a$  and  $R \sim \ell$  to the asymptotic state  $X$ . If we expand the second term in powers of  $t_{11}s_{11}$ , we can identify the  $n^{\text{th}}$  term as the contribution from transmission through the region  $R \sim a$  and reflection from the region  $R \sim \ell$ , followed by  $n$  reflections from both the regions  $R \sim a$  and  $R \sim \ell$ , followed by a final reflection from the region  $R \sim a$  and transmission through the region  $R \sim \ell$  to the state  $X$ . With each reflection from the region  $R \sim \ell$ , the amplitude decreases by a factor  $e^{-2\eta_*}$  due to the loss of probability into high-energy atom-molecule scattering states.

To calculate the dimer relaxation rate into a specific high-energy atom-molecule scattering state  $X$ , we need the S-matrix element  $S_{X,AD}$  in (6). It is extremely sensitive to the details of the interaction potential at short distances through the factor  $t_{X1}$ . However, the inclusive dimer relaxation rate summed over all high-energy atom-molecule scattering states  $X$  is much less sensitive to short distances. The matrix elements  $t_{X1}$  enter only in a combination that, by the unitarity of  $t$  and the explicit expression for  $t_{11}$  in (5), depends only on  $\eta_*$ :  $\sum_X |t_{X1}|^2 = 1 - e^{-4\eta_*}$ . Squaring the S-matrix element (6) and summing over  $X$ , we obtain

$$\sum_X |S_{X,AD}|^2 = (1 - e^{-4\eta_*})|s_{12}|^2|1 - t_{11}s_{11}|^{-2}. \quad (7)$$

This radial law shows that the inclusive dimer relaxation rate depends on the interactions only through the parameters  $a$ ,  $\Lambda_*$ , and  $\eta_*$ .

To calculate the dimer relaxation rate constant  $\beta$ , we need the behavior of the matrix  $s$  as  $k \rightarrow 0$ , where  $k$  is the wave number of the atom or dimer in the center-of-mass frame. For small  $k$ , the most general form for the entries of  $s$  that is allowed by unitarity and the analyticity of  $S_{AD,AD}$  as a function of  $k^2$  is [8]

$$\begin{aligned} s_{11} &\rightarrow -e^{2i\phi'}[1 - 2(b_0 + ib_2)ak + \dots], \\ s_{12} &\rightarrow e^{i\phi'}\sqrt{4b_0ak}[1 - (2b_3 + ib_1 + ib_2)ak + \dots], \\ s_{22} &\rightarrow 1 - 2(b_0 + ib_1)ak + \dots, \end{aligned} \quad (8)$$




FIG. 2: The dimer relaxation rate constant  $\beta$  in units of  $\hbar a/m$  as a function of  $a\Lambda_*$  for  $\eta_* = 0.1, 0.5, 1.0$ , and  $\infty$ .

where  $b_0$ ,  $b_1$ ,  $b_2$ ,  $b_3$ , and  $\phi'$  are real constants. Inserting these expressions for  $s_{ij}$  into (4), and expanding to first order in  $k$ , we find that the (complex) atom-dimer scattering length has the form

$$a_{AD} = (b_1 - b_0 \tan[s_0 \ln(a\Lambda_*) + \phi + i\eta_*])a. \quad (9)$$

For  $\eta_* = 0$ ,  $\text{Im } a_{AD}$  vanishes and Eq. (9) reduces to Efimov's expression for the atom-dimer scattering length in the absence of deep molecules [8]. The numerical constants  $b_1 = 1.46$ ,  $b_0 = 2.15$ , and  $\phi = 0.09$  were first calculated by Bedaque, Hammer, and van Kolck using an effective field theory for short-range interactions [9].

The low-energy dimer relaxation rate can be obtained by inserting the expansions (8) into the radial law (7). The rate constant  $\beta$  defined in (1) is given by the same numerical constants  $b_0$  and  $\phi$  that appear in (9):

$$\beta = \frac{20.3 \sinh(2\eta_*)}{\sin^2[s_0 \ln(a\Lambda_*) - 1.48] + \sinh^2 \eta_*} \frac{\hbar a}{m}. \quad (10)$$

Alternatively, we can use unitarity to obtain  $\beta$  directly from (9):  $\beta = -(6\pi\hbar/m) \text{Im } a_{AD}$ . In Fig. 2, we show the dimer relaxation rate constant  $\beta$  as a function of  $a\Lambda_*$  for a range of  $\eta_*$ . For small  $\eta_*$ ,  $\beta$  is strongly enhanced at the values of  $a$  corresponding to an Efimov resonance near the atom-dimer threshold. At the same values of  $a$ , there is resonant enhancement of the threshold cross section  $4\pi|a_{AD}|^2$  for atom-dimer scattering. As  $\eta_*$  increases, the resonance structure is washed out and the coefficient of  $\hbar a/m$  in (10) approaches a constant:  $\beta \approx 40.5 \hbar a/m$ .

For completeness, we also give the universal expressions for the 3-body recombination rate constants when there are deep molecules. Denoting the event rate constant by  $\alpha$ , the loss rate of low-energy atoms from a cold gas is  $dn_A/dt = -3\alpha n_A^3$ . In a BEC, the coefficient of  $n_A^3$  is smaller by a factor of 6 [20]. If  $a < 0$ , 3-body recombination can only produce deep molecules, while if  $a > 0$  it can also produce the shallow dimer. In Ref. [12], the rate

constant  $\alpha$  for  $a > 0$  was calculated as a function of  $a$  and  $\Lambda_*$  if deep molecules are absent. The effects of deep molecules on this contribution to  $\alpha$  can be deduced using the methods described above. In Ref. [14], the contribution to  $\alpha$  from recombination into deep molecules was calculated as a function of  $a$  and  $\Lambda_*$  for both signs of  $a$  and infinitesimal  $\eta_*$ . The methods described above can be used to generalize those results to arbitrary  $\eta_*$ . The two contributions to the rate constant for  $a > 0$  are

$$\alpha_{\text{shallow}} = 16.8 \frac{\hbar a^4}{m} (1 + e^{-4\eta_*} - 2e^{-2\eta_*} \cos[2s_0 \ln(a\Lambda_*) + 0.38]), \quad (11)$$

$$\alpha_{\text{deep}} = 16.8 \frac{\hbar a^4}{m} (1 - e^{-4\eta_*}). \quad (12)$$

The only contribution to  $\alpha$  for  $a < 0$  is

$$\alpha_{\text{deep}} = \frac{4590 \sinh(2\eta_*)}{\sin^2[s_0 \ln(|a|\Lambda_*) - 1.38] + \sinh^2 \eta_*} \frac{\hbar a^4}{m}. \quad (13)$$

If we set  $\eta_* = 0$  in (11), we recover the expression derived in Ref. [12]. If we expand (12, 13) to first order in  $\eta_*$ , we recover the expressions in Ref. [14] after the identification  $\eta_* = 0.2h_1$  and correcting for a missing factor of  $32\pi^2$  in Ref. [14]. The expression in [14] for  $\alpha_{\text{deep}}$  in the case  $a > 0$  has unphysical divergences that are artifacts of the expansion to first order in  $\eta_*$ . In the general expression (13), the divergences are replaced by resonance peaks.

There have been recent measurements of 3-body recombination rates for atoms with large scattering length. In Ref. [4], the rate constant  $\alpha = \frac{1}{3}K_3$  was measured for  $^{87}\text{Rb}$  atoms in the  $|1, +1\rangle$  hyperfine state in the region of the Feshbach resonance near 1007 G. It increases by at least two orders of magnitude at the resonance. Since the off-resonant scattering length is not large, the expressions (11)–(13) apply only in the resonance region and there were not enough data points in this region to observe the scaling behavior  $\alpha \sim a^4$ . In Ref. [5], the rate constant  $\alpha$  was measured for  $^{133}\text{Cs}$  atoms in the  $|3, +3\rangle$  hyperfine state in the interval  $10 \text{ G} < B < 150 \text{ G}$ , which includes several Feshbach resonances. Since the off-resonant scattering length for  $^{133}\text{Cs}$  is large, the expressions (11)–(13) should apply except near the points where  $a$  goes through zero. The data of Ref. [5] provides beautiful confirmation of the scaling behavior  $\alpha \sim a^4$ . No evidence of oscillatory dependence of the coefficient of  $a^4$  on  $\ln(|a|)$  was observed, but neither was it excluded.

Some recent theoretical predictions of 3-body recombination rates near a Feshbach resonance have given results inconsistent with the scaling behavior  $\alpha \sim a^4$ . In Ref. [21], the scaling behavior  $\alpha_{\text{deep}} \sim a^2$  was obtained by assuming a two-step process involving resonant production of the shallow dimer followed by dimer relaxation into deep molecules. Such a two-step process requires a weak-coupling assumption that breaks down in the strong-coupling region where  $a$  is large. In Ref. [22], the authors claim that  $\alpha$  scales as  $a^2$  for  $a > 0$  and as  $|a|^3$  for  $a < 0$  in a particular model for a Feshbach resonance.

However their approximations were too crude to extract the true asymptotic scaling behavior.

The rate constants  $\alpha$  and  $\beta$  in (10)–(13) can be used in mean-field descriptions of condensates of atoms and dimers near a Feshbach resonance to take into account the loss of atoms through 3-body recombination and dimer relaxation. While the scattering length  $a$  diverges as the magnetic field  $B$  passes through the Feshbach resonance, the 3-body parameters  $\Lambda_*$  and  $\eta_*$  are smooth functions of  $B$  that can be approximated by constants if

the resonance is sufficiently narrow [14]. The study of loss processes from condensates could be used to determine the parameters  $\Lambda_*$  and  $\eta_*$  for a given Feshbach resonance. Once these parameters are known, the binding energies and widths of the Efimov states can be predicted and one can develop a strategy for discovering these remarkable states.

This research was supported by DOE grant DE-FG02-91-ER4069. We thank G. Shlyapnikov for discussions.

- 
- [1] E. Tiesinga, B.J. Verhaar, and H.T.C. Stoof, Phys. Rev. A **47**, 4114 (1993); A.J. Moerdijk, B.J. Verhaar, and A. Axelsson, Phys. Rev. A **51**, 4852 (1995).
  - [2] S. Inouye *et al.*, Nature **392**, 151 (1998); J. Stenger *et al.*, Phys. Rev. Lett. **82**, 2422 (1999).
  - [3] J.L. Roberts, N.R. Claussen, S.L. Cornish, and C.E. Wieman, Phys. Rev. Lett. **85**, 728 (2000).
  - [4] A. Marte *et al.*, Phys. Rev. Lett. **89**, 283202 (2002).
  - [5] T. Weber *et al.*, Phys. Rev. Lett. **91**, 123201 (2003).
  - [6] N.R. Claussen, E.A. Donley, S.T. Thompson, and C.E. Wieman, Phys. Rev. Lett. **89**, 010401 (2002).
  - [7] E.A. Donley, N.R. Claussen, S.T. Thompson, and C.E. Wieman, Nature **417**, 529 (2002).
  - [8] V.N. Efimov, Sov. J. Nucl. Phys. **12**, 589 (1971); **29**, 546 (1979).
  - [9] P.F. Bedaque, H.-W. Hammer, and U. van Kolck, Phys. Rev. Lett. **82**, 463 (1999); Nucl. Phys. A **646**, 444 (1999).
  - [10] E. Nielsen and J.H. Macek, Phys. Rev. Lett. **83**, 1566 (1999);
  - [11] B.D. Esry, C.H. Greene, and J.P. Burke, Phys. Rev. Lett. **83**, 1751 (1999).
  - [12] P.F. Bedaque, E. Braaten, and H.-W. Hammer, Phys. Rev. Lett. **85**, 908 (2000).
  - [13] F.M. Pen'kov, Phys. Rev. A **60**, 3756 (1999); E. Nielsen, H. Suno, and B.D. Esry, Phys. Rev. A **66**, 012705 (2002).
  - [14] E. Braaten and H.-W. Hammer, Phys. Rev. Lett. **87**, 160407 (2001).
  - [15] F.A. van Abeelen and B.J. Verhaar, Phys. Rev. Lett. **83**, 1550 (1999); V.A. Yurovsky, A. Ben-Reuven, P.S. Julienne, and C.J. Williams, Phys. Rev. A **60**, R765 (1999).
  - [16] E. Braaten, H.-W. Hammer, and M. Kusunoki, Phys. Rev. A **67**, 022505 (2003).
  - [17] See e.g., S.K. Adhikari, Phys. Rev. A **66**, 013611 (2002).
  - [18] G. T. Bodwin, E. Braaten, and G. P. Lepage, Phys. Rev. D **51**, 1125 (1995) [Erratum-ibid. D **55**, 5853 (1997)].
  - [19] E. Nielsen, D.V. Fedorov, A.S. Jensen, and E. Garrido, Phys. Rep. **347**, 373 (2001).
  - [20] Yu. Kagan, B.V. Svistunov, and G.V. Shlyapnikov, JETP Lett. **42**, 209 (1985).
  - [21] V.A. Yurovsky and A. Ben-Reuven, Phys. Rev. A **67**, 050701(R) (2003).
  - [22] O.I. Kartavtsev and J.H. Macek, Few-Body Syst. **31**, 249 (2002).